

ENSO: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
2 May 2016

Outline

Summary

Recent Evolution and Current Conditions

Oceanic Niño Index (ONI)

Pacific SST Outlook

U.S. Seasonal Precipitation and Temperature Outlooks

Summary

Summary

ENSO Alert System Status: El Niño Advisory/ La Niña Watch

El Niño is present and is weakening.*

Positive equatorial sea surface temperature (SST) anomalies continue across most of the Pacific Ocean.

A transition to ENSO-neutral is likely during late Northern Hemisphere spring or early summer 2016, with an increasing chance of La Niña during the second half of the year.*


* Note: These statements are updated once a month (2nd Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).

Recent Evolution of Equatorial Pacific SST Departures (°C)

From June 2015 through September 2015, the largest positive sea surface temperature (SST) anomalies shifted westward.

Since January 2016, SST anomalies have decreased in the eastern equatorial Pacific Ocean.



Since mid-April, below-average SSTs are apparent near S. America, while above-average SSTs remain in the central and east-central Pacific.



Niño Region SST Departures (°C) Recent Evolution


The latest weekly SST departures are:

Niño 4	0.9°C
Niño 3.4	0.8°C
Niño 3	0.4°C
Niño 1+2	-0.3°C



SST Departures ($^{\circ}\text{C}$) in the Tropical Pacific During the Last Four Weeks


During the last four weeks, equatorial SSTs were above average across the Pacific, except near S. America.



Weekly SST Departures during the Last Four Weeks


During the last four weeks, positive SST anomalies decreased in strength and negative anomalies have emerged in the eastern Pacific.

Weekly SST Anomalies (DEG C)



Change in Weekly SST Departures over the Last Four Weeks

During the last four weeks, equatorial SST anomalies significantly decreased across the eastern Pacific.




Upper-Ocean Conditions in the Equatorial Pacific

The basin-wide equatorial upper ocean (0-300 m) heat content is greatest prior to and during the early stages of a Pacific warm (El Niño) episode (compare top 2 panels), and least prior to and during the early stages of a cold (La Niña) episode.

The slope of the oceanic thermocline is least (greatest) during warm (cold) episodes.


Recent values of the upper-ocean heat anomalies (near zero) and thermocline slope index (negative) reflect a weakening El Niño.

The monthly thermocline slope index represents the difference in anomalous depth of the 20°C isotherm between the western Pacific (160°E-150°W) and the eastern Pacific (90°-140°W).




Central and Eastern Pacific Upper-Ocean (0-300 m) Weekly Average Temperature Anomalies

Positive anomalies decreased during November and December, increased during the first half of January 2016, and have significantly decreased since late January with negative values appearing during March/April.




Sub-Surface Temperature Departures in the Equatorial Pacific

During the last two months, the vertical extent of the layer of positive subsurface temperature anomalies in the central and eastern equatorial Pacific has steadily decreased.



Below-average subsurface anomalies are evident across most of the equatorial Pacific and extend to the surface in the eastern Pacific.




Tropical OLR and Wind Anomalies During the Last 30 Days

Negative OLR anomalies (enhanced convection and precipitation) were evident over the central Pacific. Positive OLR anomalies (suppressed convection and precipitation) were observed over the Philippines, Malaysia, and northern Australia.

Low-level (850-hPa) winds were near average across the equatorial Pacific.

Anomalous upper-level (200-hPa) easterlies were observed over the western and central equatorial Pacific. Anomalous anti-cyclones straddled the equator.



Intraseasonal Variability

Intraseasonal variability in the atmosphere (wind and pressure), which is often related to the Madden-Julian Oscillation (MJO), can significantly impact surface and subsurface conditions across the Pacific Ocean.

Related to this activity:


Significant weakening of the low-level easterly winds usually initiates an eastward-propagating oceanic Kelvin wave.

Weekly Heat Content Evolution in the Equatorial Pacific

Downwelling phases of a Kelvin wave were observed in mid-May to late June, July-August, and October to November, and January-February 2016.

Since February 2016, an upwelling phase of a Kelvin wave has influenced the equatorial Pacific.

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Down-welling and warming occur in the leading portion of a Kelvin wave, and up-welling and cooling occur in the trailing portion.




Low-level (850-hPa) Zonal (east-west) Wind Anomalies (m s^{-1})

During early January and late February 2016 westerly wind bursts were observed between 140°E and 140°W .

Recently, winds were mostly near average across the equatorial Pacific.

Westerly Wind Anomalies (orange/red shading)

Easterly Wind Anomalies (blue shading)




Upper-level (200-hPa) Velocity Potential Anomalies

Throughout the period, anomalous upper-level divergence (green shading) and convergence (brown shading) have generally persisted over the Central/Eastern Pacific and Indonesia, respectively.

Sub-seasonal or Madden-Julian Oscillation (MJO) activity contributed to an eastward propagation of regions of upper-level divergence and convergence during December 2015-January 2016 and February-March 2016.


Unfavorable for precipitation (brown shading)
Favorable for precipitation (green shading)



Outgoing Longwave Radiation (OLR) Anomalies

Throughout the period negative anomalies have been observed over the central Pacific, and positive anomalies have persisted near Indonesia or the western Pacific.

Drier-than-average Conditions (orange/red shading)
Wetter-than-average Conditions (blue shading)



Oceanic Niño Index (ONI)

The ONI is based on SST departures from average in the Niño 3.4 region, and is a principal measure for monitoring, assessing, and predicting ENSO.

Defined as the three-month running-mean SST departures in the Niño 3.4 region. Departures are based on a set of improved homogeneous historical SST analyses (Extended Reconstructed SST - ERSST.v4). The SST reconstruction methodology is described in Huang et al., 2015, J. Climate, vol. 28, 911-930.)

It is one index that helps to place current events into a historical perspective

NOAA Operational Definitions for El Niño and La Niña

El Niño: characterized by a positive ONI greater than or equal to $+0.5^{\circ}\text{C}$.

La Niña: characterized by a negative ONI less than or equal to -0.5°C .


By historical standards, to be classified as a full-fledged El Niño or La Niña episode, these thresholds must be exceeded for a period of at least 5 consecutive overlapping 3-month seasons.

CPC considers El Niño or La Niña conditions to occur when the monthly Niño3.4 OISST departures meet or exceed $\pm 0.5^{\circ}\text{C}$ along with consistent atmospheric features. These anomalies must also be forecasted to persist for 3 consecutive months.

ONI (°C): Evolution since 1950

The most recent ONI value (January - March 2016) is 2.0°C.


El Niño ↑
Neutral
La Niña ↓



CPC/IRI Probabilistic ENSO Outlook

Updated: 14 April 2016

The chance of El Niño gradually decreases during the spring and ENSO-neutral is favored during May-June-July (MJJ) 2016. The chance of La Niña and ENSO-neutral are about equal by June-July-August (JJA) 2016, with La Niña favored afterwards.



IRI/CPC Pacific Niño 3.4 SST Model Outlook

Positive Niño 3.4 SST anomalies are predicted to weaken through 2016.

Most models suggest a transition to ENSO-neutral by May-June-July (MJJ) 2016 with the possibility of La Niña conditions during the Northern Hemisphere fall and winter.






Figure provided by the International Research Institute (IRI) for Climate and Society (updated 19 April 2016).

SST Outlook: NCEP CFS.v2 Forecast (PDF corrected)

Issued: 2 May 2016


The CFS.v2 ensemble mean (black dashed line) predicts Neutral by May-June-July (MJJ) and La Niña after June-July-August (JJA) 2016.



Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

From mid March to mid April, the overall pattern was characterized mostly by anomalous ridging (and above-average temperatures) over western N. America and an anomalous trough (and below-average temperatures) over portions of eastern N. America.


During the last half of April, an anomalous ridge (and above-average temperatures) dominated Alaska, western Canada, and much of the contiguous U.S.



Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

From mid March to mid April, the overall pattern was characterized mostly by anomalous ridging (and above-average temperatures) over western N. America and an anomalous trough (and below-average temperatures) over portions of eastern N. America.


During the last half of April, an anomalous ridge (and above-average temperatures) dominated Alaska, western Canada, and much of the contiguous U.S.



Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

From mid March to mid April, the overall pattern was characterized mostly by anomalous ridging (and above-average temperatures) over western N. America and an anomalous trough (and below-average temperatures) over portions of eastern N. America.


During the last half of April, an anomalous ridge (and above-average temperatures) dominated Alaska, western Canada, and much of the contiguous U.S.




U.S. Temperature and Precipitation Departures During the Last 30 Days

End Date: 30 April 2016

Percent of Average Precipitation




Temperature Departures (degree C)




U.S. Temperature and Precipitation Departures During the Last 90 Days

End Date: 30 April 2016

Percent of Average Precipitation





Temperature Departures (degree C)



U. S. Seasonal Outlooks

May - July 2016

The seasonal outlooks combine the effects of long-term trends, soil moisture, and, when appropriate, ENSO.



Summary

ENSO Alert System Status: El Niño Advisory/La Niña Watch

El Niño is present and is weakening.*

Positive equatorial sea surface temperature (SST) anomalies continue across most of the Pacific Ocean.

A transition to ENSO-neutral is likely during late Northern Hemisphere spring or early summer 2016, with an increasing chance of La Niña during the second half of the year.*

* Note: These statements are updated once a month (2nd Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).