

# ENSO: Recent Evolution, Current Status and Predictions



Update prepared by:  
Climate Prediction Center / NCEP  
1 August 2016

# Outline

Summary

Recent Evolution and Current Conditions

Oceanic Niño Index (ONI)

Pacific SST Outlook

U.S. Seasonal Precipitation and Temperature Outlooks

Summary

# Summary

## ENSO Alert System Status: La Niña Watch

ENSO-neutral conditions are present.\*

Equatorial sea surface temperatures (SST) are near or below average in the east-central and eastern Pacific Ocean.


La Niña is favored to develop during August-October (ASO)2016, with about a 55-60% chance of La Niña during the fall and winter 2016-17.\*

\* Note: These statements are updated once a month (2<sup>nd</sup> Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).

# Recent Evolution of Equatorial Pacific SST Departures (°C)

Since January 2016, SST anomalies have decreased in the eastern equatorial Pacific Ocean.



Since mid-April, near-to-below average SSTs are apparent in the eastern Pacific, while above-average SSTs remain west of the Date Line.



# Niño Region SST Departures (°C) Recent Evolution


The latest weekly SST departures are:

Niño 4	0.2°C
Niño 3.4	-0.5°C
Niño 3	-0.6°C
Niño 1+2	0.1°C




# SST Departures (°C) in the Tropical Pacific During the Last Four Weeks

During the last four weeks, equatorial SSTs were above average west of the International Date Line, and near-to-below average across the east-central and eastern Pacific.



# Global SST Departures (°C) During the Last Four Weeks


During the last four weeks, tropical SSTs were above average in the western Pacific, the eastern Indian Ocean, and the Atlantic. Equatorial SSTs were near or below average in the east-central and eastern Pacific Ocean and the western Indian Ocean.



# Weekly SST Departures during the Last Four Weeks

During the last four weeks, negative anomalies persisted in the east-central and eastern Pacific.


Weekly SST Anomalies (DEG C)





# Change in Weekly SST Departures over the Last Four Weeks

During the last four weeks, equatorial SST anomalies mostly decreased across the eastern Pacific Ocean.




# Upper-Ocean Conditions in the Equatorial Pacific

The basin-wide equatorial upper ocean (0-300 m) heat content is greatest prior to and during the early stages of a Pacific warm (El Niño) episode (compare top 2 panels), and least prior to and during the early stages of a cold (La Niña) episode.

The slope of the oceanic thermocline is least (greatest) during warm (cold) episodes.


Recent values of the upper-ocean heat anomalies (negative) and thermocline slope index (near zero) reflect ENSO-neutral conditions.

*The monthly thermocline slope index represents the difference in anomalous depth of the 20°C isotherm between the western Pacific (160°E-150°W) and the eastern Pacific (90°-140°W).*




# Central and Eastern Pacific Upper-Ocean (0-300 m) Weekly Average Temperature Anomalies

Positive anomalies decreased during November and December, increased during the first half of January 2016, and then decreased significantly from late January through early May with negative values evident since early March. Since late May, the negative anomalies have weakened.




# Sub-Surface Temperature Departures in the Equatorial Pacific

During the last two months, below-average subsurface temperatures were evident across the equatorial Pacific, extending to the surface in the east-central and eastern Pacific.



Recently, the below-average temperatures have weakened across the eastern Pacific Ocean.






# Tropical OLR and Wind Anomalies During the Last 30 Days

Negative OLR anomalies (enhanced convection and precipitation) were evident over parts of Indonesia. Regions of positive OLR anomalies (suppressed convection and precipitation) were observed around the International Date Line.

Low-level (850-hPa) winds were near average across the equatorial Pacific.

Upper-level (200-hPa) winds were anomalously easterly over a small region of the east-central equatorial Pacific.



# Intraseasonal Variability

Intraseasonal variability in the atmosphere (wind and pressure), which is often related to the Madden-Julian Oscillation (MJO), can significantly impact surface and subsurface conditions across the Pacific Ocean.

Related to this activity:

Significant weakening of the low-level easterly winds usually initiates an eastward-propagating oceanic Kelvin wave.


# Weekly Heat Content Evolution in the Equatorial Pacific

Downwelling phases of equatorial oceanic Kelvin waves were observed in October to November and January-February 2016.

Since the passage of an upwelling equatorial oceanic Kelvin wave in March 2016, below-average subsurface temperatures have continued across much of the equatorial Pacific.

Since June 2016, the below-average subsurface temperatures have weakened across the equatorial Pacific.

Equatorial oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and upwelling and cooling occur in the trailing portion.





# Low-level (850-hPa) Zonal (east-west) Wind Anomalies ( $\text{m s}^{-1}$ )


During late February 2016, a westerly wind burst was observed between  $140^\circ\text{E}$  and  $140^\circ\text{W}$ .

Since February, the equatorial Pacific has been characterized by weak anomalous westerlies and easterlies.

In the last week, weak wind anomalies were evident across most of the equatorial Pacific.

Westerly Wind Anomalies (orange/red shading)

Easterly Wind Anomalies (blue shading)






# Upper-level (200-hPa) Velocity Potential Anomalies

Through April 2016, anomalous upper-level divergence (green shading) and convergence (brown shading) generally persisted over the central/eastern Pacific and Indonesia, respectively.

Sub-seasonal or Madden-Julian Oscillation (MJO) activity contributed to an eastward propagation of regions of upper-level divergence and convergence during February-March and from mid April through the present.

Unfavorable for precipitation (brown shading)  
Favorable for precipitation (green shading)




# Outgoing Longwave Radiation (OLR) Anomalies

Through April, negative OLR anomalies were observed over the central Pacific, and positive anomalies persisted near Indonesia or the western Pacific.

Currently, negative OLR anomalies predominate near Indonesia.

Drier-than-average Conditions (orange/red shading)  
Wetter-than-average Conditions (blue shading)



# Oceanic Niño Index (ONI)

The ONI is based on SST departures from average in the Niño 3.4 region, and is a principal measure for monitoring, assessing, and predicting ENSO.

Defined as the three-month running-mean SST departures in the Niño 3.4 region. Departures are based on a set of improved homogeneous historical SST analyses (Extended Reconstructed SST - ERSST.v4). The SST reconstruction methodology is described in Huang et al., 2015, J. Climate, vol. 28, 911-930.)

It is one index that helps to place current events into a historical perspective

# NOAA Operational Definitions for El Niño and La Niña

El Niño: characterized by a positive ONI greater than or equal to  $+0.5^{\circ}\text{C}$ .

La Niña: characterized by a negative ONI less than or equal to  $-0.5^{\circ}\text{C}$ .


By historical standards, to be classified as a full-fledged El Niño or La Niña episode, these thresholds must be exceeded for a period of at least 5 consecutive overlapping 3-month seasons.

CPC considers El Niño or La Niña conditions to occur when the monthly Niño3.4 OISST departures meet or exceed  $\pm 0.5^{\circ}\text{C}$  along with consistent atmospheric features. These anomalies must also be forecasted to persist for 3 consecutive months.

# ONI (°C): Evolution since 1950

The most recent ONI value (April-June 2016) is 0.7°C.

El Niño ↑  
Neutral  
La Niña ↓



# Historical El Niño and La Niña Episodes Based on the ONI computed using ERSST.v4

Recent Pacific warm (red) and cold (blue) periods based on a threshold of  $\pm 0.5$  °C for the Oceanic Nino Index (ONI) [3 month running mean of ERSST.v4 SST anomalies in the Nino 3.4 region (5N-5S, 120-170W)]. For historical purposes, periods of below and above normal SSTs are colored in blue and red when the threshold is met for a minimum of 5 consecutive over-lapping seasons.


The ONI is one measure of the El Niño-Southern Oscillation, and other indices can confirm whether features consistent with a coupled ocean-atmosphere phenomenon accompanied these periods. The complete table going back to DJF 1950 can be found [here](#).

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
2004	0.3	0.2	0.1	0.1	0.2	0.3	0.5	0.7	0.7	0.7	0.7	0.7
2005	0.6	0.6	0.5	0.5	0.4	0.2	0.1	0.0	0.0	-0.1	-0.4	-0.7
2006	-0.7	-0.6	-0.4	-0.2	0.0	0.1	0.2	0.3	0.5	0.8	0.9	1.0
2007	0.7	0.3	0.0	-0.1	-0.2	-0.2	-0.3	-0.6	-0.8	-1.1	-1.2	-1.3
2008	-1.4	-1.3	-1.1	-0.9	-0.7	-0.5	-0.3	-0.2	-0.2	-0.3	-0.5	-0.7
2009	-0.8	-0.7	-0.4	-0.1	0.2	0.4	0.5	0.6	0.7	1.0	1.2	1.3
2010	1.3	1.1	0.8	0.5	0.0	-0.4	-0.8	-1.1	-1.3	-1.4	-1.3	-1.4
2011	-1.3	-1.1	-0.8	-0.6	-0.3	-0.2	-0.3	-0.5	-0.7	-0.9	-0.9	-0.8
2012	-0.7	-0.6	-0.5	-0.4	-0.3	-0.1	0.1	0.3	0.4	0.4	0.2	-0.2
2013	-0.4	-0.5	-0.3	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.2	-0.3
2014	-0.5	-0.6	-0.4	-0.2	0.0	0.0	0.0	0.0	0.2	0.4	0.6	0.6
2015	0.5	0.4	0.5	0.7	0.9	1.0	1.2	1.5	1.8	2.1	2.2	2.3
2016	2.2	1.9	1.5	1.1	0.7							

# CPC/IRI Probabilistic ENSO Outlook

Updated: 14 July 2016

La Niña is slightly favored by August-September-October (ASO) 2016. The chance of La Niña is roughly 55-60% during the Northern Hemisphere fall and winter 2016-17.



# IRI/CPC Pacific Niño

## 3.4 SST Model Outlook

Most multi-model averages indicate a borderline or a weak La Niña starting during the Northern Hemisphere fall and persisting through winter 2016-17.






Figure provided by the International Research Institute (IRI) for Climate and Society (updated 18 July 2016).



# SST Outlook: NCEP CFS.v2 Forecast (PDF corrected)


Issued: 1 August 2016

The CFS.v2 ensemble mean (black dashed line) predict ENSO-neutral during the Northern Hemisphere fall and winter 2016-17.




# Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

Since mid-June, above-average heights and temperatures have prevailed over the southern tier of the U.S.




# Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

Since mid-June, above-average heights and temperatures have prevailed over the southern tier of the U.S.



# Atmospheric anomalies over the North Pacific and North America During the Last 60 Days


Since mid-June, above-average heights and temperatures have prevailed over the southern tier of the U.S.




# U.S. Temperature and Precipitation Departures During the Last 30 Days

End Date: 30 July 2016

Percent of Average Precipitation



Temperature Departures (degree C)







# U.S. Temperature and Precipitation Departures During the Last 90 Days

End Date: 30 July 2016

Percent of Average Precipitation



Temperature Departures (degree C)




# U. S. Seasonal Outlooks


August - October 2016

The seasonal outlooks combine the effects of long-term trends, soil moisture, and, when appropriate, ENSO.

## Precipitation



## Temperature



# Summary

## ENSO Alert System Status: La Niña Watch

ENSO-neutral conditions are present.\*

Equatorial sea surface temperatures (SST) are near or below average in the east-central and eastern Pacific Ocean.

La Niña is favored to develop during August-October (ASO)2016, with about a 55-60% chance of La Niña during the fall and winter 2016-17.\*

\* Note: These statements are updated once a month (2<sup>nd</sup> Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).