

ENSO: Recent Evolution, Current Status and Predictions



Update prepared by:
Climate Prediction Center / NCEP
29 May 2017

Outline

Summary

Recent Evolution and Current Conditions

Oceanic Niño Index (ONI)

Pacific SST Outlook

U.S. Seasonal Precipitation and Temperature Outlooks

Summary

Summary

ENSO Alert System Status: Not Active

ENSO-neutral conditions are present.*

Equatorial sea surface temperatures (SSTs) are near-to-above average across most of the Pacific Ocean.

ENSO-neutral and El Niño are nearly equally favored during the Northern Hemisphere summer and fall 2017.*


* Note: These statements are updated once a month (2nd Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).

Recent Evolution of Equatorial Pacific SST Departures (°C)

From July through December 2016, below average SSTs were observed over most of the central and eastern Pacific Ocean.

During January and February 2017, above-average SSTs expanded within the eastern Pacific Ocean.



Since mid April 2017, near-to-above average SSTs expanded across the equatorial Pacific.



Niño Region SST Departures (°C) Recent Evolution


The latest weekly SST departures are:

| | |
|----------|-------|
| Niño 4 | 0.3°C |
| Niño 3.4 | 0.6°C |
| Niño 3 | 0.6°C |
| Niño 1+2 | 0.4°C |




SST Departures (°C) in the Tropical Pacific During the Last Four Weeks

During the last four weeks, equatorial SSTs were near-average near the International Date Line, and above-average across the far western Pacific and portions of the eastern Pacific.



Global SST Departures (°C) During the Last Four Weeks


During the last four weeks, equatorial SSTs were above average in the Atlantic, western Indian, and the eastern and western Pacific Oceans, and were near-average in the central Pacific Ocean.



Weekly SST Departures during the Last Four Weeks


During the last four weeks, near-average SSTs continued in the central Pacific, while above-average SSTs weakened across the eastern Pacific.

Weekly SST Anomalies (DEG C)



Change in Weekly SST Departures over the Last Four Weeks

During the last four weeks, negative changes in equatorial SST anomalies were evident in regions of the eastern Pacific.






Upper-Ocean Conditions in the Equatorial Pacific

The basin-wide equatorial upper ocean (0-300 m) heat content is greatest prior to and during the early stages of a Pacific warm (El Niño) episode (compare top 2 panels), and least prior to and during the early stages of a cold (La Niña) episode.

The slope of the oceanic thermocline is least (greatest) during warm (cold) episodes.


Recent values of the upper-ocean heat anomalies (near average) and thermocline slope index (near average) reflect ENSO-neutral conditions.

The monthly thermocline slope index represents the difference in anomalous depth of the 20°C isotherm between the western Pacific (160°E-150°W) and the eastern Pacific (90°-140°W).




Central and Eastern Pacific Upper-Ocean (0-300 m) Weekly Average Temperature Anomalies

Negative subsurface temperature anomalies were present through December 2016. Positive anomalies were present from mid-January through March 2017 before weakening to near zero. From mid-April to mid-May, positive anomalies strengthened. Recently, positive anomalies have weakened slightly.




Sub-Surface Temperature Departures in the Equatorial Pacific

During the last two months, positive subsurface temperature anomalies have shifted from the western to the central Pacific.



Recently, positive anomalies have weakened near the Date Line.




Tropical OLR and Wind Anomalies During the Last 30 Days

Negative OLR anomalies (enhanced convection and precipitation) were evident across portions of Indonesia and south of the equator across the western and central Pacific. Positive OLR anomalies (suppressed convection and precipitation) were observed north of the equator in the central Pacific.

Low-level (850-hPa) winds were anomalously westerly in the far eastern equatorial Pacific and were anomalous easterly in a small region of the western Pacific.

Upper-level (200-hPa) easterly wind anomalies were evident over the eastern Pacific.



Intraseasonal Variability

Intraseasonal variability in the atmosphere (wind and pressure), which is often related to the Madden-Julian Oscillation (MJO), can significantly impact surface and subsurface conditions across the Pacific Ocean.

Related to this activity:

Significant weakening of the low-level easterly winds usually initiates an eastward-propagating oceanic Kelvin wave.


Weekly Heat Content Evolution in the Equatorial Pacific

From March 2016 through December 2016, below-average subsurface temperatures extended across most of the equatorial Pacific.

From February 2017 through April 2017, positive subsurface temperature anomalies persisted in the western and eastern Pacific Ocean, with oceanic Kelvin waves resulting in anomalous temperature variability in the central Pacific.

Since early May 2017, positive subsurface temperature anomalies shifted eastward into the central and east-central Pacific.

Equatorial oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Down-welling and warming occur in the leading portion of a Kelvin wave, and up-welling and cooling occur in the trailing portion.



Low-level (850-hPa) Zonal (east-west) Wind Anomalies (m s^{-1})

From mid September 2016 to mid April 2017, low-level easterly wind anomalies generally persisted over the central and western equatorial Pacific.


Since January 2017, westerly wind anomalies were generally observed over the eastern Pacific Ocean.

During mid-to-late April 2017, low-level westerly anomalies expanded to the western and central Pacific.

Since mid May, easterly wind anomalies returned to the west-central Pacific, while westerly anomalies persisted in the eastern Pacific.

Westerly Wind Anomalies (orange/red shading)

Easterly Wind Anomalies (blue shading)




Upper-level (200-hPa) Velocity Potential Anomalies

From September 2016 to present, anomalous upper-level divergence (green shading) generally persisted near Indonesia.

Eastward propagation of regions of upper-level divergence (green shading) and convergence (brown shading) is particularly evident during November 2016, January-February 2017, and since early April 2017.

Unfavorable for precipitation (brown shading)
Favorable for precipitation (green shading)

Note: Eastward propagation is not necessarily indicative of the Madden-Julian Oscillation (MJO).




Outgoing Longwave Radiation (OLR) Anomalies

From early September 2016 to mid April 2017, positive OLR anomalies persisted near the International Date Line, with negative OLR anomalies persisting near the Maritime Continent/far western Pacific Ocean.

In the last week, OLR anomalies were negative across a small portion of Indonesia and the Indian Ocean.

Drier-than-average Conditions (orange/red shading)
Wetter-than-average Conditions (blue shading)



Oceanic Niño Index (ONI)

The ONI is based on SST departures from average in the Niño 3.4 region, and is a principal measure for monitoring, assessing, and predicting ENSO.

Defined as the three-month running-mean SST departures in the Niño 3.4 region. Departures are based on a set of improved homogeneous historical SST analyses (Extended Reconstructed SST - ERSST.v4). The SST reconstruction methodology is described in Huang et al., 2015, J. Climate, vol. 28, 911-930.)

It is one index that helps to place current events into a historical perspective

NOAA Operational Definitions for El Niño and La Niña

El Niño: characterized by a positive ONI greater than or equal to $+0.5^{\circ}\text{C}$.

La Niña: characterized by a negative ONI less than or equal to -0.5°C .


By historical standards, to be classified as a full-fledged El Niño or La Niña episode, these thresholds must be exceeded for a period of at least 5 consecutive overlapping 3-month seasons.

CPC considers El Niño or La Niña conditions to occur when the monthly Niño3.4 OISST departures meet or exceed $\pm 0.5^{\circ}\text{C}$ along with consistent atmospheric features. These anomalies must also be forecasted to persist for 3 consecutive months.

ONI (°C): Evolution since 1950

The most recent ONI value (February - April 2017) is 0.1°C.


El Niño ↑
Neutral
La Niña ↓



CPC/IRI Probabilistic ENSO Outlook

Updated: 11 May 2017

ENSO-neutral is favored through spring 2017, with nearly equal chances (~45%) of El Niño and ENSO-neutral through the remainder of 2017.



IRI/CPC Pacific Niño

3.4 SST Model Outlook

Many models favor at least a weak El Niño by the Northern Hemisphere summer 2017, continuing through winter 2017-18.





Figure provided by the International Research Institute (IRI) for Climate and Society (updated 16 May 2017).

SST Outlook: NCEP CFS.v2 Forecast (PDF corrected)

Issued: 29 May 2017


The CFS.v2 ensemble mean (black dashed line) favors ENSO-neutral to continue through 2017.

CFSv2 forecast Nino3.4 SST anomalies (K) (PDF corrected)



— Latest 8 forecast members
— Earliest 8 forecast members
— Other forecast members
- - - Forecast ensemble mean
— NCDP daily analysis


(Model bias correct base period: 1999–2010; Climatology base period: 1982–2010)



Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

The pattern of temperature, winds, and heights have been variable over the last 60 days.


More recently, an anomalous trough and negative temperature anomalies have been evident over the central U.S.



Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

The pattern of temperature, winds, and heights have been variable over the last 60 days.


More recently, an anomalous trough and negative temperature anomalies have been evident over the central U.S.



Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

The pattern of temperature, winds, and heights have been variable over the last 60 days.


More recently, an anomalous trough and negative temperature anomalies have been evident over the central U.S.




U.S. Temperature and Precipitation Departures During the Last 30 Days

End Date: 27 May 2017

Percent of Average Precipitation




Temperature Departures (degree C)




U.S. Temperature and Precipitation Departures During the Last 90 Days

End Date: 27 May 2017

Percent of Average Precipitation



Temperature Departures (degree C)




U. S. Seasonal Outlooks


June - August 2017

The seasonal outlooks combine the effects of long-term trends, soil moisture, and, when appropriate, ENSO.

Precipitation



Temperature



Summary

ENSO Alert System Status: Not Active

ENSO-neutral conditions are present.*

Equatorial sea surface temperatures (SSTs) are near-to-above average across most of the Pacific Ocean.

ENSO-neutral and El Niño are nearly equally favored during the Northern Hemisphere summer and fall 2017.*

* Note: These statements are updated once a month (2nd Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).